

ON FOX COLORINGS OF SYMMETRIC UNIONS

TOSHIFUMI TANAKA

Abstract. A symmetric union in the 3-space \mathbb{R}^3 is a knot, obtained from a knot in \mathbb{R}^3 and its mirror image, which are symmetric with respect to a 2-plane in \mathbb{R}^3 , by taking the connected sum of them and moreover by connecting them with some vertical twists along the plane. In this paper, we study the p -coloring of a symmetric union.

Key words: Symmetric union; p -coloring; partial knot.

1. INTRODUCTION

A *symmetric union* was introduced by Kinoshita and Terasaka [2]. A symmetric union is known to be a *ribbon knot* which bounds a smooth disk in the 4-ball. (See [5] for the definition.) Every ribbon knot with minimal crossing number ≤ 10 is a symmetric union [1, 3] and it is known that all two-bridge ribbon knots can be represented as symmetric unions. (See [4, 6].)

A p -coloring of a diagram D of a knot is an assignment of one of the numbers $0, 1, \dots, p-1$ to each arc of D such a way that at each crossing the sum of the numbers of the under-crossings is equal to twice the number of the over-crossing modulo p . The number of p -colorings of D is a knot invariant. We denote it by $c_p(K)$ for a knot K . A p -coloring is said to be *non-trivial* if it has at least two numbers. A knot K is p -colorable if K has a diagram which admits a non-trivial p -coloring.

In this paper, we study p -colorings of a symmetric union and show the following theorem.

Theorem 1.1. *Let \hat{K} be a knot with a symmetric union presentation with a partial knot K and p , a positive odd prime integer. Then K is p -colorable if and only if \hat{K} is p -colorable.*

Corollary 1.2. *Let \hat{K} be a knot with a symmetric union presentation with a partial knot K and p , a positive odd prime integer. Then $c_p(K) = p$ if and only if $c_p(\hat{K}) = p$.*

Acknowledgements. The author is partially supported by the Ministry of Education, Science, Sports and Culture, Grant-in-Aid for Scientific Research(C), 2019-2021 (19K03465).

2. THE DEFINITION OF A SYMMETRIC UNION

Let \mathbb{R}^3 be the 3-space with x -, y -, and z -axes. Let $\mathbb{R}_+^3 = \{(x, y, z) \in \mathbb{R}^3 \mid x > 0\}$ and $\mathbb{R}_-^3 = \{(x, y, z) \in \mathbb{R}^3 \mid x < 0\}$. Throughout this paper, a *tangle* denotes two arcs properly embedded in a 3-ball. We denote the tangle made of $|m|$ half-twists along the z -axis as a diagram by an integer $m \in \mathbb{Z}$ and the horizontal trivial tangle by ∞ with respect to the x -axis as in Figure 1. A symmetric union is defined as follows.

Definition 2.1. We take a knot \tilde{K} in \mathbb{R}_-^3 and its mirror image \tilde{K}^* in \mathbb{R}_+^3 such that \tilde{K} and \tilde{K}^* are symmetric with respect to the yz -plane \mathbb{R}_{yz}^2 as in Figure 2(a). Here we consider a diagram of a knot in the xz -plane \mathbb{R}_{xz}^2 and we denote the diagrams of \tilde{K} and \tilde{K}^* by \tilde{D} and \tilde{D}^* , respectively. We take 0-tangles T_0, T_1, \dots, T_k as in Figure 2(a). Then we replace the tangles T_0, T_1, \dots, T_k with tangles $\infty, m_1, m_2, \dots, m_k$ as in Figure 2(b). (See Figure 3 for example.) Here we assume that $m_i \neq \infty$ ($1 \leq i \leq k$). The resultant diagram is called a *symmetric union presentation* and we denote it by $\tilde{D} \cup \tilde{D}^*(m_1, \dots, m_k)$. The knot \tilde{K} is called the *partial knot* of the symmetric union.

2010 Mathematics Subject Classification. Primary 57M25; Secondary 57M27.

If a knot K has a diagram $\tilde{D} \cup \tilde{D}^*(m_1, \dots, m_k)$, then the knot K is called a *symmetric union*.




FIGURE 1




FIGURE 2




FIGURE 3

3. PROOFS

Proof of Theorem 1.1. By a result of [3], we know that $\det(\hat{K}) = (\det(K))^2$. Note that $\det(\hat{K})$ is divisible by p if and only if $\det(K)$ is divisible by p . On the other hand, it is known that a knot is p -colorable if and only if p divides its determinant when p is an odd prime integer. [7] Thus we know that \hat{K} is p -colorable if and only if K is p -colorable.

4. THE NUMBER OF COLORINGS OF A KNOT

We can easily show the following result since we can obtain a p -coloring of a symmetric union from a p -coloring of the partial knot.

Proposition 4.1. *Let \hat{K} be a knot with a symmetric union presentation with a partial knot K and p , a positive integer. Then $c_p(K) \leq c_p(\hat{K})$.*

Proposition 4.2. [8] *We have $c_p(K_1)c_p(K_2) = p \cdot c_p(\hat{K})$ if \hat{K} is the connected sum of K_1 and K_2 .*

Now we consider the following question.

Question. Let \hat{K} be a knot with a symmetric union presentation with a partial knot K . Does the equality $c_p(K)^2 = p \cdot c_p(\hat{K})$ always hold?

Example 4.3. We take a p -coloring of 6_1 as in Figure 5.




FIGURE 4

Then we have the following congruences:

$$2x_1 \equiv x_3 + x_4 \pmod{p}$$

$$2x_4 \equiv x_6 + x_1 \pmod{p}$$

$$2x_6 \equiv x_1 + x_2 \pmod{p}$$

$$2x_2 \equiv x_6 + x_5 \pmod{p}$$

$$2x_5 \equiv x_2 + x_3 \pmod{p}$$

$$2x_3 \equiv x_4 + x_5 \pmod{p}$$

Thus we have the following congruences:

$$-x_3 + 6x_5 - 5x_6 \equiv 0 \pmod{p}$$

$$-x_1 + 4x_5 - 3x_6 \equiv 0 \pmod{p}$$

$$-x_2 - 4x_5 + 5x_6 \equiv 0 \pmod{p}$$

$$-9x_5 + 9x_6 \equiv 0 \pmod{p}$$

$$-x_4 + 2x_5 - x_6 \equiv 0 \pmod{p}$$

$$9x_5 - 9x_6 \equiv 0 \pmod{p}$$

In the case when $p \neq 3m$, we have $x_5 \equiv x_6$ by the fourth congruence. Thus we have $x_1 \equiv x_2 \equiv x_3 \equiv x_4 \equiv x_5 \equiv x_6$ by the other congruences. Therefore we know that $c_p(6_1) = p$.

In the case when $p = 3m$ and $m \neq 3n$, we have $x_5 \equiv x_6 \pmod{m}$ by the fourth congruence. Then we have $9m$ choices for the pair (x_5, x_6) . Thus we know that $c_p(6_1) = 9m = 3p$.

In the case when $p = 3m$ and $m = 3n$, we have $x_5 \equiv x_6 \pmod{n}$ by the fourth congruence. Then we have $81n$ choices for the pair (x_5, x_6) . Thus we know that $c_p(6_1) = 81n = 9p$.

Thus we have

$$(1) \quad c_p(6_1) = \begin{cases} 9p & (p = 9n \ (n \in \mathbb{N})) \\ 3p & (p = 3m, m \neq 3n \ (m, n \in \mathbb{N})) \\ p & (p \neq 3m \ (m \in \mathbb{N})) \end{cases} .$$

On the other hand, 6_1 has a symmetric union presentation with 3_1 as the partial knot as follows:




FIGURE 5

We can easily calculate $c_p(3_1)$ as follows:

$$(2) \quad c_p(3_1) = \begin{cases} 3p & (p = 3m \ (m \in \mathbb{N})) \\ p & (p \neq 3m \ (m \in \mathbb{N})) \end{cases} .$$

So we know that $c_p(3_1)^2 \neq p \cdot c_p(6_1)$ if $p = 3m, m \neq 3n \ (m, n \in \mathbb{N})$. By using Proposition 4.2, we can obtain infinitely many symmetric unions with such a property.

REFERENCES

1. M. Eisermann and C. Lamm: *Equivalence of symmetric union diagrams*, J. Knot Theory Ramifications **16** (2007), no. 7, 879–898.
2. S. Kinoshita and H. Terasaka: *On unions of knots*, Osaka J. Math. Vol. 9 (1957), 131–153.
3. C. Lamm: *Symmetric unions and ribbon knots*, Osaka J. Math., Vol. 37 (2000), 537–550.
4. C. Lamm: *Symmetric union presentations for 2-bridge ribbon knots*, arxiv:math.GT/0602395, 2006.
5. W. B. R. Lickorish: *An introduction to knot theory*, Graduate Texts in Mathematics, **175**, Springer-Verlag, New York, 1997.
6. P. Lisca: *Lens spaces, rational balls and the ribbon conjecture*, Geom. Topol. **11** (2007), 429–472.
7. C. Livingston: *Knot theory*, Carus Mathematical Monographs, 24. Mathematical Association of America, Washington, DC, 1993.
8. J. H. Przytycki: *3-coloring and other elementary invariants of knots*, Knot theory (Warsaw, 1995), 275–295, Banach Center Publ., 42, Polish Acad. Sci. Inst. Math., Warsaw, 1998.

DEPARTMENT OF MATHEMATICS, FACULTY OF EDUCATION, GIFU UNIVERSITY, YANAGIDO 1-1, GIFU, 501-1193, JAPAN.

Email address: tanakat@gifu-u.ac.jp