

A formula for the Jones polynomial of symmetric unions

Toshifumi Tanaka

Keyword: symmetric union, Jones polynomial, ribbon knot.

Institution: Department of Mathematics, Faculty of Education, Gifu University, Yanagido 1-1, Gifu, 501-1193, Japan.

Abstract. Symmetric unions are links obtained from a knot in the 3-space \mathbb{R}^3 by taking diagrams of the knot and its mirror image, which are symmetrically placed with respect to an axis in the 2-plane, and connecting them to obtain a diagram by inserting a 2-tangle with no twists and a finite number of 2-tangles with twists along the axis in the diagram. The Jones polynomial of a symmetric union has some interesting properties. In this paper, we give a formula of the Jones polynomial of a symmetric union.

1 Introduction

A *symmetric union* was first introduced by Kinoshita and Terasaka [4]. In recent years, Lamm [5] generalized the definition and investigated the relationship between a symmetric union and a ribbon knot. It is easy to see that every symmetric union represents a ribbon knot, on the other hand, the converse question is still open. In this paper, we give the following formula for the Jones polynomial [2] of a knot with a symmetric union presentation. Let $\bar{V}_L(t) = V_L(t)/(-t^{-1/2} - t^{1/2})^{c-1}$ for an oriented link L , where c is the number of the components of L and $V_L(t)$ is the Jones polynomial of L . (See Section 2 for the definition.)

Theorem 1.1. *Let \bar{K} be a knot with a symmetric union presentation $D_K \cup D_K^*(\infty, m_1, \dots, m_k)_1$. Then*

$$V_{\bar{K}}(t) = \sum_{(\ell_1, \dots, \ell_k)} \prod_{i=1}^k (-t^{-1})^{m_i p_i} (1 - (-t^{-1})^{m_i})^{q_i} \bar{V}_{D_K \cup D_K^*(\infty, \ell_1, \dots, \ell_k)_w}(t),$$

where $\ell_i \in \{0, \infty\}$,

$$p_i = \begin{cases} 1 & (\ell_i = 0) \\ 0 & (\ell_i = \infty) \end{cases}$$

, $q_i = 1 - p_i$ and $w = \sum_{i=1}^k q_i + 1$ for each k -tuple of tangles (ℓ_1, \dots, ℓ_k) . The sum is taken over all combinations of k tangles 0's and ∞ 's to tangles ℓ_1, \dots, ℓ_k .

The following corollary generalizes a result of Lamm in the case when $s = 1$ in [5].

Corollary 1.2. *Let \bar{K} be a link with a symmetric union presentation $D_K \cup D_K^*(\infty, m_1, \dots, m_k)_1$ and s , a positive integer. If $m_i \equiv 0 \pmod{s}$ for each i , then $V_{\bar{K}}(-\exp(\frac{2\pi i}{s})) = V_K(-\exp(\frac{2\pi i}{s})) \cdot V_{K^*}(-\exp(\frac{2\pi i}{s}))$.*

A knot is called *amphicheiral*, if it is isotopic to its mirror image. By Theorem 1.1, we have the following as special cases.

Theorem 1.3. [7] *Let \bar{K} be a knot with a symmetric union presentation $D_K \cup D_K^*(\infty, m)_1$. Then $t^m V_{\bar{K}}(t) + (-1)^m V_{\bar{K}}(t^{-1}) = (t^m + (-1)^m) V_K(t) \cdot V_K(t^{-1})$. In particular, if \bar{K} is amphicheiral, then $V_{\bar{K}}(t) = V_K(t) V_K(t^{-1})$.*

In this paper, all knots and links are oriented unless otherwise stated. In Section 2, we give the definitions of the Jones polynomial of a link and a symmetric union. In Section 3, we shall prove Theorem 1.1, Corollary 1.2 and Theorem 1.3. In Section 4, we shall consider an amphicheiral symmetric union and the property.

Acknowledgements. The author is partially supported by the Ministry of Education, Science, Sports and Culture, Grant-in-Aid for Scientific Research(C), 2019-2021 (19K03465).

2 Definitions

Definition 2.1. Let K be a link in \mathbb{R}^3 . We denote a diagram of K by D_K . The *bracket polynomial* of a diagram of K [3][6], $\langle D_K \rangle$ can be defined as a polynomial which satisfies the following identities.

- i) $\langle \bigcirc \rangle = 1$,
- ii) $\langle D_K \cup \bigcirc \rangle = -(A^2 + A^{-2}) \langle D_K \rangle$,
- iii) $\langle \begin{array}{c} \diagup \\ \diagdown \end{array} \rangle = A \langle \begin{array}{c} \diagdown \\ \diagup \end{array} \rangle + A^{-1} \langle \begin{array}{c} \diagup \\ \diagup \end{array} \rangle \langle \begin{array}{c} \diagdown \\ \diagdown \end{array} \rangle$.

We define $V_{D_K}(t) \in \mathbb{Z}[t^{1/2}, t^{-1/2}]$ by $V_{D_K}(t) = \{(-A^3)^{-\delta(D_K)} \langle D_K \rangle\}_{t^{1/2}=A^{-2}}$ for any diagram D_K for K , where $\delta(D_K)$ is the writhe of D_K . It is shown that $V_{D_K}(t)$ is an invariant of K [3]. Then we denote $V_{D_K}(t)$ by $V_K(t)$ and call it the *Jones polynomial* of K .

Here we give the definition of a symmetric union as follows. (See [5] for the original definition.) We denote the tangle made of half twists by an integer $m \in \mathbb{Z}$ and the horizontal trivial tangle by ∞ as in Fig. 1.

Definition 2.2. Let D be an unoriented diagram of a knot and D^* the diagram D reflected at an axis in the plane. We replace the tangles $T_i = 0$ ($i = 0, \dots, k$) on the symmetry axis by either $T_i = \infty$ or $T_i = m_i \in \mathbb{Z}$ for each i , providing that at least one of T_i 's is replaced by ∞ as shown in Fig. 1. We call the result a *symmetric union* and denote it by $D \cup D^*(m_0, \dots, m_k)_\mu$, where μ is the number of ∞ -tangles on the axis.




Figure 1: A symmetric union

If a link L has a diagram $D \cup D^*(m_0, \dots, m_k)_\mu$, then the diagram is called a *symmetric union presentation* for L . We say that a link L is a *symmetric union* if L has a symmetric union presentation.

3 A formula of the Jones polynomial

Let $T_m = \frac{(-A^{-3})^m - A^m}{-A^{-2} - A^2}$ and $S_m = A^m$ for an indeterminate A and a non-zero integer m .

Proof of Theorem 1.1. By using the identities of the bracket polynomial in Definition 2.1, we have
 $\langle D_K \cup D_K^*(\infty, m_1, m_2, \dots, m_k)_1 \rangle = A^{m_1/|m_1|} \langle D_K \cup D_K^*(\infty, m_1 - 1, m_2, \dots, m_k)_1 \rangle$
 $+ F_1 \langle D_K \cup D_K^*(\infty, \infty, m_2, \dots, m_k)_2 \rangle = \dots = A^{\sum_{i=1}^{|m_1|} m_1/|m_1|} \langle D_K \cup D_K^*(\infty, 0, m_2, \dots, m_k)_1 \rangle$
 $+ (F_1 + \dots + F_{|m_1|}) \langle D_K \cup D_K^*(\infty, \infty, m_2, \dots, m_k)_2 \rangle = A^{m_1} \langle D_K \cup D_K^*(\infty, 0, m_2, \dots, m_k)_1 \rangle$
 $+ (\sum_{j=1}^{|m_1|} F_j) \langle D_K \cup D_K^*(\infty, \infty, m_2, \dots, m_k)_2 \rangle$, where each F_j is some polynomial.

Now we calculate a formula of $\sum_{j=1}^{|m_1|} F_j$ by considering the unknot instead of K as follows. We assume that D_K is a diagram as in Fig. 2 so that we have a symmetric union of the unknot. We denote the diagram in Fig. 2 by D_\circ .




Figure 2: D_\circ

Then by calculating each term by using the identities in Definition 2.1, we have
 $\langle D_\circ \cup D_\circ^*(\infty, m_1, m_2, \dots, m_k)_1 \rangle = (-A^{-3m_1/|m_1|})^{|m_1|} \dots (-A^{-3m_k/|m_k|})^{|m_k|} (-A^{-2} - A^2)$
 $= (-1)^{\sum_{i=1}^k |m_i|} A^{-3 \sum_{i=1}^k m_i} (-A^{-2} - A^2)$,
 $\langle D_\circ \cup D_\circ^*(\infty, 0, m_2, \dots, m_k)_1 \rangle = (-1)^{\sum_{i=2}^k |m_i|} A^{-3 \sum_{i=2}^k m_i} (-A^{-2} - A^2)$ and
 $\langle D_\circ \cup D_\circ^*(\infty, \infty, m_2, \dots, m_k)_2 \rangle = (-1)^{\sum_{i=2}^k |m_i|} A^{-3 \sum_{i=2}^k m_i} (-A^{-2} - A^2)^2$.

Then we have

$$\begin{aligned} \sum_{j=1}^{|m_1|} F_j &= \frac{\langle D_\circ \cup D_\circ^*(\infty, m_1, m_2, \dots, m_k)_1 \rangle - \langle D_\circ \cup D_\circ^*(\infty, 0, m_2, \dots, m_k)_1 \rangle}{\langle D_\circ \cup D_\circ^*(\infty, \infty, m_2, \dots, m_k)_2 \rangle} \\ &= \frac{(-1)^{\sum_{i=1}^k |m_i|} A^{-3 \sum_{i=1}^k m_i} (-A^{-2} - A^2) - (-1)^{\sum_{i=2}^k |m_i|} A^{-3 \sum_{i=2}^k m_i} (-A^{-2} - A^2)}{(-1)^{\sum_{i=2}^k |m_i|} A^{-3 \sum_{i=2}^k m_i} (-A^{-2} - A^2)^2} \\ &= \frac{(-1)^{|m_1|} A^{-3m_1} - A^{m_1}}{-A^{-2} - A^2} = \frac{(-A^{-3})^{m_1} - A^{m_1}}{-A^{-2} - A^2}. \end{aligned}$$

Thus

$$\begin{aligned} \langle D_K \cup D_K^*(\infty, m_1, m_2, \dots, m_k)_1 \rangle &= A^{m_1} \langle D_K \cup D_K^*(\infty, 0, m_2, \dots, m_k)_1 \rangle \\ &+ \frac{(-A^{-3})^{m_1} - A^{m_1}}{-A^{-2} - A^2} \langle D_K \cup D_K^*(\infty, \infty, m_2, \dots, m_k)_2 \rangle \\ &= S_{m_1} \langle D_K \cup D_K^*(\infty, 0, m_2, \dots, m_k)_1 \rangle + T_{m_1} \langle D_K \cup D_K^*(\infty, \infty, m_2, \dots, m_k)_2 \rangle. \end{aligned}$$

By iterating the same argument, we have

$$\begin{aligned} \langle D_K \cup D_K^*(\infty, m_1, m_2, \dots, m_k)_1 \rangle &= S_{m_1} S_{m_2} \langle D_K \cup D_K^*(\infty, 0, 0, m_3, \dots, m_k)_1 \rangle \\ &+ S_{m_1} T_{m_2} \langle D_K \cup D_K^*(\infty, 0, \infty, m_3, \dots, m_k)_2 \rangle \\ &+ T_{m_1} S_{m_2} \langle D_K \cup D_K^*(\infty, \infty, 0, m_3, \dots, m_k)_2 \rangle \end{aligned}$$

$$+T_{m_1}T_{m_2} \langle D_K \cup D_K^*(\infty, \infty, \infty, m_3, \dots, m_k)_3 \rangle.$$

Then by induction on k , we know that

$$\begin{aligned} \langle D_K \cup D_K^*(\infty, m_1, m_2, \dots, m_k)_1 \rangle &= \sum_{(\ell_1, \dots, \ell_k)} \prod_{i=1}^k \Pi_{m_i}^{p_i(\ell_1, \dots, \ell_k)} T_{m_i}^{q_i(\infty, \ell_1, \dots, \ell_k)} \times \\ \langle D_K \cup D_K^*(\infty, \ell_1, \ell_2, \dots, \ell_k)_w \rangle, & \text{ where } \ell_i \in \{0, \infty\}, p_i(\ell_1, \dots, \ell_k) = 1 \text{ or } 0 \text{ if } \ell_i = 0 \text{ or } \infty \text{ respectively,} \\ q_i(\ell_1, \dots, \ell_k) &= 1 - p_i(\ell_1, \dots, \ell_k) \text{ and } w = \sum_{i=1}^k q_i(\ell_1, \dots, \ell_k) + 1. \end{aligned}$$

Now we obtain that

$$\begin{aligned} V_{D_K \cup D_K^*(\infty, m_1, \dots, m_k)_1}(A) &= (-A)^{-3 \sum_{i=1}^k (-m_i)} \langle D_K \cup D_K^*(\infty, m_1, m_2, \dots, m_k)_1 \rangle \\ &= (-A^3)^{\sum_{i=1}^k m_i} \langle D_K \cup D_K^*(\infty, m_1, m_2, \dots, m_k)_1 \rangle = (-A^3)^{\sum_{i=1}^k m_i} \times \\ &\sum_{(\ell_1, \dots, \ell_k)} \prod_{i=1}^k \Pi_{m_i}^{p_i(\ell_1, \dots, \ell_k)} \left(\frac{(-A^{-3})^{m_i} - A^{m_i}}{-A^{-2} - A^2} \right)^{q_i(\ell_1, \dots, \ell_k)} \langle D_K \cup D_K^*(\infty, \ell_1, \dots, \ell_k)_w \rangle \\ &= \sum_{(\ell_1, \dots, \ell_k)} \prod_{i=1}^k (-1)^{m_i p_i(\ell_1, \dots, \ell_k)} A^{4m_i p_i(\ell_1, \dots, \ell_k)} \left(\frac{1 - (-1)^{m_i} A^{4m_i}}{-A^{-2} - A^2} \right)^{q_i(\ell_1, \dots, \ell_k)} \langle D_K \cup D_K^*(\infty, \ell_1, \dots, \ell_k)_w \rangle. \end{aligned}$$

Using $t^{1/2} = A^{-2}$, we have

$$\begin{aligned} V_{D_K \cup D_K^*(\infty, m_1, \dots, m_k)_1}(t) &= \sum_{(\ell_1, \dots, \ell_k)} \prod_{i=1}^k (-1)^{m_i p_i(\ell_1, \dots, \ell_k)} t^{-m_i p_i(\ell_1, \dots, \ell_k)} \left(\frac{1 - (-1)^{m_i} t^{-m_i}}{-t^{1/2} - t^{-1/2}} \right)^{q_i(\ell_1, \dots, \ell_k)} V_{D_K \cup D_K^*(\infty, \ell_1, \dots, \ell_k)_w}(t) \\ &= \sum_{(\ell_1, \dots, \ell_k)} \prod_{i=1}^k (-t^{-1})^{m_i p_i(\ell_1, \dots, \ell_k)} \left(\frac{1 - (-1)^{m_i} t^{-m_i}}{-t^{1/2} - t^{-1/2}} \right)^{q_i(\ell_1, \dots, \ell_k)} V_{D_K \cup D_K^*(\infty, \ell_1, \dots, \ell_k)_w}(t). \end{aligned}$$

Therefore we obtain

$$V_{D_K \cup D_K^*(\infty, m_1, \dots, m_k)_1}(t) = \sum_{(\ell_1, \dots, \ell_k)} \prod_{i=1}^k (-t^{-1})^{m_i p_i(\ell_1, \dots, \ell_k)} (1 - (-1)^{m_i} t^{-m_i})^{q_i(\ell_1, \dots, \ell_k)} \bar{V}_{D_K \cup D_K^*(\infty, \ell_1, \dots, \ell_k)_w}(t).$$

Remark 3.1. By a result of Eisermann ([1], Theorem 1), we know that $\bar{V}_{D_K \cup D_K^*(\infty, \ell_1, \dots, \ell_k)_w}(t)$ in the statement of Theorem 1.1 is always a Laurent polynomial. The integers p_i , q_i and w depend on k -tuple of tangles (ℓ_1, \dots, ℓ_k) , however we abbreviate the integers for simplicity in the statement of Theorem 1.1.

Proof of Corollary 1.2. By Theorem 1.1, we have $V_{\bar{K}}(-\exp(\frac{2\pi i}{s})) = \bar{V}_{D_K \cup D_K^*(\infty, 0, \dots, 0)_1}(-\exp(\frac{2\pi i}{s})) = V_K(-\exp(\frac{2\pi i}{s})) \cdot V_{K^*}(-\exp(\frac{2\pi i}{s}))$.

Proof of Theorem 1.3. In the case when $k = 1$ in Theorem 1.1, we have the formula as follows.

$$V_{\bar{K}}(t) = (-1)^m t^{-m} \bar{V}_{D_K \cup D_K^*(\infty, 0)_1}(t) + (1 - (-1)^m t^{-m}) \bar{V}_{D_K \cup D_K^*(\infty, \infty)_2}(t).$$

Then the first part of the theorem is obtained as follows.

$$\begin{aligned} t^m V_{\bar{K}}(t) + (-1)^m V_{\bar{K}}(t^{-1}) &= \{(-1)^m \bar{V}_{D_K \cup D_K^*(\infty, 0)_1}(t) + (t^m - (-1)^m) \bar{V}_{D_K \cup D_K^*(\infty, \infty)_2}(t)\} + \\ &\{t^m \bar{V}_{D_K \cup D_K^*(\infty, 0)_1}(t) + ((-1)^m - t^m) \bar{V}_{D_K \cup D_K^*(\infty, \infty)_2}(t)\} \\ &= (t^m + (-1)^m) \bar{V}_{D_K \cup D_K^*(\infty, 0)_1}(t) = (t^m + (-1)^m) V_K(t) V_K(t^{-1}). \end{aligned}$$

The latter part of the theorem follows immediately from the first part because $V_{\bar{K}}(t) = V_{\bar{K}}(t^{-1})$ if \bar{K} is amphicheiral [6].

4 Examples

For a positive integer m , let K_m be a knot with the symmetric union presentation as described in Fig. 3. It is easily seen that K_m is an amphicheiral knot for each m .

If $m = 1$, then the Jones polynomial of K_m is calculated by Theorem 1.1 as follows.




Figure 3: K_m

$$V_{K_1}(t) = -t^{-5} + 3t^{-4} - 7t^{-3} + 10t^{-2} - 12t^{-1} + 15 - 12t + 10t^2 - 7t^3 + 3t^4 - t^5.$$

By a calculation in [7], we know that $V_t(K_m)$ cannot have a form $f(t)f(t^{-1})$ for any Laurent polynomial $f(t) \in \mathbb{Z}[t, t^{-1}]$. Thus, by Theorem 1.3, we know that K_1 cannot have a symmetric union presentation $D_K \cup D_K^*(\infty, m)_1$. The Jones polynomials of K_2 and K_3 are calculated as follows.

$$V_{K_2}(t) = t^{-7} - 2t^{-6} + 2t^{-5} + t^{-4} - 7t^{-3} + 12t^{-2} - 16t^{-1} + 19 - 16t + 12t^2 - 7t^3 + t^4 + 2t^5 - 2t^6 + t^7$$

$$V_{K_3}(t) = -t^{-8} + 2t^{-7} - 3t^{-6} + t^{-5} + 5t^{-4} - 14t^{-3} + 24t^{-2} - 31t^{-1} + 35 - 31t + 24t^2 - 14t^3 + 5t^4 + t^5 - 3t^6 + 2t^7 - t^8$$

By using Mathematica [8], we know that $V_{K_2}(t)$ and $V_{K_3}(t)$ are irreducible. In particular, they cannot have a form $f(t)f(t^{-1})$ for any Laurent polynomial $f(t)$.

Question. Does $V_{K_m}(t)$ have a form $f(t)f(t^{-1})$ for some Laurent polynomial $f(t) \in \mathbb{Z}[t, t^{-1}]$ for each $m > 1$?

References

- [1] M. Eisermann, The Jones polynomial of ribbon links. *Geom. Topol.* 13 (2009), no. 2, 623–660.
- [2] V. F. R. Jones, *A polynomial invariant for knots via von Neumann algebras*, *Bull. Amer. Math. Soc. (N.S.)* 12 (1985), no. 1, 103–111.
- [3] L. H. Kauffman, *State models and the Jones polynomials*, *Topology*. Vol. 26 (1987), 395–407.
- [4] S. Kinoshita and H. Terasaka, *On unions of knots*, *Osaka J. Math.* Vol. 9 (1957), 131–153.
- [5] C. Lamm, *Symmetric unions and ribbon knots*, *Osaka J. Math.*, Vol. 37 (2000), 537–550.
- [6] W. B. R. Lickorish, *An introduction to knot theory*, *Graduate Texts in Mathematics*, 175.
- [7] T. Tanaka, *The Jones polynomial of knots with symmetric union presentations*, *J. Korean Math. Soc.* **52** (2015), no. 2, 389–402.
- [8] Wolfram Research, Inc., *Mathematica*, Version 11.2. Champaign, IL, 2017.

