



岐阜大学機関リポジトリ

Gifu University Institutional Repository

## A Function of Extracellular $Ca^{2+}$ in Controls on Cell Survival and Cell Death During Chicken Embryogenesis

メタデータ	言語: English 出版者: 公開日: 2022-06-08 キーワード: 作成者: NAGANAWA, Hidetoshi, IWASAWA, Atsushi, NAKAMURA, Takao メールアドレス: 所属:
URL	<a href="http://hdl.handle.net/20.500.12099/5895">http://hdl.handle.net/20.500.12099/5895</a>

# A Function of Extracellular $\text{Ca}^{2+}$ in Controls on Cell Survival and Cell Death During Chicken Embryogenesis

Hidetoshi NAGANAWA, Atsushi IWASAWA and Takao NAKAMURA

*Department of Animal Science and Technology*

*(Received July 19, 1999)*

## SUMMARY

This paper describes a function of extracellular calcium ion ( $\text{Ca}^{2+}$ ) that may maintain normal embryonic development by means of controlling the dose level. Experimental results *in vivo* show that an excess of  $\text{Ca}^{2+}$  outside the cell administered to the chick embryo (after 24–30 hr of incubation) caused either the induction or inhibition of cell death, depending on the dose.

Res. Bull. Fac. Agr. Gifu Univ. (64): 21–25, 1999

## INTRODUCTION

Up to now, only a few investigations concerning the role of extracellular  $\text{Ca}^{2+}$  on cell death have been performed<sup>1)</sup>. In normal embryonic development, the cell death (called programmed cell death or apoptosis) occurs at specific times and sites<sup>2–13)</sup>. This process is characterized by DNA degradation into fragments the size of one or more nucleosomes (DNA ladder)<sup>14,15)</sup> and by specific morphological changes including chromatin condensation and nuclear fragmentation<sup>4,8,13)</sup>.

In the present study, we have shown that excess extracellular  $\text{Ca}^{2+}$  administered to the chick embryo caused either the induction or inhibition of cell death, depending on the dose. This dual function of extracellular  $\text{Ca}^{2+}$  on cell survival and cell death during embryogenesis suggests that controlling the level of this factor may maintain normal embryonic development. During our studies we detected the above phenomena (chromatin condensation, nuclear fragmentation) by supravital staining (see Materials and Methods).

## MATERIALS AND METHODS

Fertilized hen's eggs (Plymouth Rock), incubated at 38°C, were used throughout.

*Staging of embryo development.* The staging of embryo development was based on morphological characteristics, and was designated according to the Eguchi stage seriation (a revision of Hamburger-Hamilton's)<sup>16,17)</sup>.

*Extracellular ion modification.* Various doses of  $\text{K}^+$  or  $\text{Ca}^{2+}$  dissolved in distilled water were administered to an embryo through a window made in the egg shell via the chorioallantoic membrane (0.2 ml per embryo) at 24 hr of incubation. The sealed egg was allowed to develop further until 30 hr of incubation.

*Methods of counting dead cells.* Dead cells were distinguished from living cells by supravital staining and/or histochemical (metachromatic) methods. For supravital staining, the entire removed embryos were incubated in 0.01% (w/w) Nile blue sulfate at 37°C for 20 min, and washed twice with modified Ringer's saline solution<sup>3)</sup>. Because most dead cells are phagocytosed by macrophages (or by neighboring cells), the number of phagocytosed cells (chromatin condensation, nuclear fragmentation) can be used as an index of cell death<sup>18)</sup>. Counting the number of phagocytosed cells was performed on the photographs of stained embryos. Data were expressed as the percent of dead cells by comparing the photographs, ( $\times 400$ ) of adjacent serial sections each stained with Nile blue and hematoxylin-eosin.

## RESULTS AND DISCUSSION

Numbers of apoptotic cells as functions of the extracellular free cations are given in Figure 1. The  $x=100$  concentration corresponds to the normal value of the extracellular cation concentration of the chicken ( $[K^+]_o$ : 5.63 mM,  $[Ca^{2+}]_o$ : 1.63 mM). The apoptotic cell numbers in neuroblasts increased with increasing doses of extracellular  $K^+$  with a plateau at higher concentrations. On the other hand, the apoptotic cell numbers in neuroblasts increased with a peak at  $x=110$  concentration, then decreased with increasing doses of extracellular  $Ca^{2+}$ . This was also observed in somitic cells to a lesser extent. No apparent changes in apoptotic cell numbers were observed in hemocytoblasts or epidermoblasts.

The increase in apoptotic cell numbers with a modification of extracellular  $K^+$  might be caused by a large intracellular inflow of  $K^+$ , which could facilitate intracellular pathways leading to apoptosis. Increased intracellular  $Ca^{2+}$  could result from  $Ca^{2+}$  inflow from extracellular fluid through membrane potential-dependent  $Ca^{2+}$  channels (which might be activated by the  $K^+$  inflow). However, to induce apoptosis, cells need an increase in intracellular  $Ca^{2+}$  of about a hundred-fold over the physiological, extracellular  $Ca^{2+}$  concentration<sup>1)</sup>, and hence,  $Ca^{2+}$  inflow might not be a candidate inducer of apoptosis. Nevertheless, the induction or inhibition of the apoptotic process in neuroblastic and somitic cells depending on the dose of extracellular  $Ca^{2+}$  (as shown in Figure 1) suggests that the role of extracellular cations in embryos is different from that in adult animals. Moreover, it is probable that the effect of extracellular  $Ca^{2+}$  on apoptosis differs between cell types.

To conclude, we found in this study that extracellular  $Ca^{2+}$  as a control on cell survival or cell death is feasible, and that such a cation could contribute to the delicate developmental regulation during apoptosis. To confirm our assumptions about extracellular cation activities, we need to measure the intracellular ion concentrations and membrane potential of the cells at various stages during apoptosis.

## ACKNOWLEDGMENTS

This paper is part of a master's thesis submitted to the Graduate School of Agriculture, Gifu University by H. N. We thank the committee members, Prof. Michiharu Kamiyoshi and Dr. Osamu Doi, for their helpful discussions. Special thanks also to Ms. Tatiana I. Orgiljanova for her assistance.




Fig.1. Apoptosis induced by excess extracellular cations(K<sup>+</sup>, Ca<sup>2+</sup>). Various doses of these cations were administered to a chick embryo at 24 hr of incubation. Dead cells were counted 6 hr later.

## REFERENCES

- 1) Naganawa, H.: Programmed Cell Death (Apoptosis) in Teratogenesis of the Chick Embryo and the Effective Teratogens on Ontogeny. 212 pp. (Master's thesis, Graduate School of Agriculture, Gifu University), 1998. [In Japanese]
- 2) Glücksmann, A.: Cell deaths in normal vertebrate ontogeny. *Biol. Revs. Camb. Philos. Soc.* 26: 59-86, 1951.
- 3) Saunders, J. W., Jr.: Death in embryonic systems: death in cells in the usual accompaniment of embryonic growth and differentiation. *Science* 154: 604-612, 1966.
- 4) Kerr, J. R. F., Wyllie, A. H., & Currie, A. R.: Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *Br. J. Cancer* 26: 239-257, 1972.
- 5) Saunders, J. W., Jr., Gasseling, M. T., & Saunders, L. C.: Cellular death in morphogenesis of the avian wing. *Develop. Biol.* 5: 147-178, 1962.
- 6) Wyllie, A. H.: Apoptosis: cell death under homeostatic control. *Arch. Toxicol.* 11: 3-10, 1987.
- 7) Wyllie, A. H.: Apoptosis: cell death in tissue regulation. *J. Pathol.* 153: 313-316, 1987.
- 8) Gerschenson, L. E., & Rotello, R. J.: Apoptosis: a different type of cell death. *FASEB J.* 6: 2450-2455, 1992.
- 9) Ellis, R. E., Yuan, J., & Horvitz, H. R.: Mechanisms and functions of cell death. *Annu. Rev. Cell Biol.* 7: 663-698, 1991.
- 10) Alison, M. R., & Sarraf, C. E.: Apoptosis: a gene-directed programme of cell death. *J. R. Coll. Physicians Lond.* 26: 25-35, 1992.
- 11) Cotter, T. G., Lennon, S. V., Glynn, J. G., & Martin, S. J.: Cell death via apoptosis and its relationships to growth, development and differentiation of both tumour and normal cells. *Anticancer Res.* 10: 1153-1160, 1990.
- 12) Kerr, J. R. F., Searle, J., Hamon, B. V., & Bishop, C. J.: 'Apoptosis', in "Perspectives in Mammalian Cell Death" Potten, C. S. ed. New York: Oxford University Press, 93-128, 1987.
- 13) Searle, J., Kerr, J. F. R., & Bishop, C. J.: Necrosis and apoptosis: distinct modes of cell death with fundamentally different significance. *Pathol. Annu.* 17: 229-259, 1982.
- 14) Bartner, C. D., Oldenburg, N. B. E., & Cidlowski, J. A.: The role of DNA fragmentation in apoptosis. *Trends Cell Biol.* 5: 21-26, 1995.
- 15) Eastman, A., & Barry, M. A.: The origins of DNA breaks: a consequence of DNA damage, DNA repair, or apoptosis? *Cancer Invest.* 10: 229-240, 1992.
- 16) Eguchi, G.: 'Developmental stage seriation of the chick embryo', in "Experiments for Developmental Biology" Ishihara, K. ed. Tokyo: Kyoritsu Publishing Co., 253-259, 1980. [In Japanese]
- 17) Hamburger, V., & Hamilton, H. L.: A series of normal stages in the development of the chick embryo. *J. Morphol.* 88: 49-92, 1951.
- 18) Toné, S., Tanaka, S., & Kato, Y.: The inhibitory effect of 5-Bromodeoxyuridine of the programmed cell death in the chick limb. *Develop., Growth and Differ.* 25: 381-391, 1983.

# 胚発生過程の細胞生存および細胞死調節における細胞外カルシウムイオンの作用

長縄秀俊・岩澤 淳・中村孝雄

動物生産学講座

(1998年7月19日受理)

## 要 約

本論文は細胞外カルシウムイオンのレベルが正常な胚発生に及ぼす作用について検討した。インビボにおける試験結果は、鶏胚に孵卵24時間目から30時間目の間に投与した細胞外のカルシウムイオンが、その投与量によって細胞死の誘導も阻害も引き起こすことを示した。

岐阜大農研報 (64) : 21-25, 1999